Triumph HealthCare Presents Debricare in Wound Care



Indications:

Debridement of necrotic tissues & liquefecation of slough in acute and chronic lesions, like

- Pressure Ulcers
- Burns
- Post-operative wounds
- Pilondial cyst wounds
- Carbuncles
- Traumatic & Infected Wounds

"Papain, the proteolytic enzyme from the fruit of carica papaya, is a potent digestant of non-viable protien matter but is harmless to viable tissue. It is active over a pH range of 3 to 12. Papain is relatively ineffective when used alone as a debriding agent and requires the presence of activators to stimulate its digestive potency. in Debricare (Papain and Urea), Papain is combined with Urea, a denaturant of protiens, to bring about two supplimental chemical actions: (1) to expose by solvent action the activators of Papain, and (2) to denature the non-viable protien matter in lesions and thereby render it more susceptible to enzymatic digestion. Pharmacological studies have shown that the combination of Papain and Urea result as much digestive activity as Papain alone"

Debricare

Debricare

Debricare

Debricare

15 gm Tube @ Rs.139/-



9-167/2/13/1/A. TGS Nilayam, Boduppal, Hyderabad-92

Triumph HealthCare Presents Debricare in Wound Care



DESCRIPTION:

Debricare (Papain and Urea) enzymatic Debriding ointment contains Papain IP greater than or equal to 521700 units of activity and Urea IP 100 mg in a hydrophilic ointment base.

INDICATIONS:

Debricare (Papain and Urea) is indicated for Debridement of necrotic and liquefaction of slough in acute and chronic lesions such as pressure ulcers, varicose and diabetic ulcars, burns, post-poperative wounds, pilondial cyst wounds, carbuncles and miscellaneous traumatic or infected wounds.

DOSAGE AND ADMINISTRATION:

Cleanse the wound with wound cleanser or saline. Avoid cleansing with hydrogen peroxide solution as it may inactivate the Papain. Apply **Debricare** (Papain and Urea) directly to the wound, cover with appropriate dressing, secure into place. Once Daily or twice daily applications are preferred. Irrigate the wound at each dressing to remove any accumulation of liquefied necrotic material.

Note: Papain may also be inactivated by the salts of heavy metals such as lead, silver and mercury. Contact with medications containing these metals should be avoided.

HOW SUPPLIED: Debricare (Papain and Urea) is available as 15gm lamitube. Store in a cool place.

SIDE EFFECTS:

Debricare (Papain and Urea) ointment is generally well tolerated and non-irritatting. A transient "burning" sensation may be experienced by a small percentage of patients upon applying Debricare (Papain and Urea) ointment. Occasionally, the profuse exudates from enzymatic digestion may irritate the skin. In such cases, more frequent dressing changes will alleviate discomfort until exudates clears.

PRECAUTIONS: Not to be used in Eyes.

CONTRAINDICATIONS:

Do not use if you are allergic to or have known or suspected hypersensitivity to any in-gradients in this product

Debricare Debricare Debricare



9-167/2/13/1/A. TGS Nilayam, Boduppal, Hyderabad-92